

Chronic Urticaria References

Akoglu, G., Atakan, N., Cakir, B., Kalayci, O., & Hayran, M. (2012). Effects of low pseudoallergen diet on urticarial activity and leukotriene levels in chronic urticaria. *Arch Dermatol Res*, 304(4), 257-262. <https://doi.org/10.1007/s00403-011-1203-3>

Anuradha, P., Maiti, R., Jyothirmai, J., Mujeebuddin, O. & Anuradha, M. (2010). Loratadine versus levocetirizine in chronic idiopathic urticaria: a comparative study of the efficacy and safety. *Indian J Pharmacol*, 42(1), 12-16. <https://doi.org/10.4103/0253-7613.62399>

Arkwright, P.D. (2009). Anti-CD20 or anti-IgE therapy for severe chronic autoimmune urticaria. *J Allergy Clin Immunol*, 123(2), 510-511. <https://doi.org/10.1016/j.jaci.2008.11.043>

Asero, R. (2007). Chronic unremitting urticaria: is the use of antihistamines above the licensed dose effective. A preliminary study of cetirizine at licensed and above-licensed doses. *Clin Exp Dermatol*, 32(1), 34-8. <https://doi.org/10.1111/j.1365-2230.2006.02278.x>

Asero, R., Tedeschi, A., & Cugno, M. (2010). Heparin and Tranexamic Acid therapy may be effective in treatment-resistant chronic urticaria with elevated d-dimer: a pilot study. *Int Arch Allergy Immunol*, 152(4), 384-389. <https://doi.org/10.1159/000292947>

Asero, R., & Tedeschi, A. (2010). Usefulness of a short course of oral prednisone in antihistamine-resistant chronic urticaria: a retrospective analysis. *J Investig Allergol Clin Immunol*, 20, 386-90. <https://pubmed.ncbi.nlm.nih.gov/20945604/>

Asero R, Tedeschi A, & Cugno M. (2013). Treatment of refractory chronic urticaria: current and future therapeutic options. *Am J Clin Dermatol*, 14(6), 481-488. <https://doi.org/10.1007/s40257-013-0047-3>

Augustin, M., & Ehrle S. (2009). Safety and efficacy of desloratadine in chronic idiopathic urticaria in clinical practice: an observational study of 9246 patients. *J Eur Acad Dermatol Venereol*, 23(3), 292-299. <https://doi.org/10.1111/j.1468-3083.2008.03034.x>

Aversano, M., Caiazzo, P., Iorio, G., Ponticiello, L., Laganá, B., & Leccese, F. (2005). Improvement of chronic idiopathic urticaria with L-thyroxine: a new TSH role in immune response? *Allergy*, 60(4), 89-93. <https://doi.org/10.1111/j.1398-9995.2005.00723.x>

- Babu, K.S., Arshad, S.H., & Holgate, S.T. (2001). Omalizumab, a novel anti-IgE therapy in allergic disorders. *Expert Opin Biol Ther*, 1(6), 1049-1058. <https://doi.org/10.1517/14712598.1.6.1049>
- Bagenstose, S.E., Levin, L., & Bernstein, J.A. (2004). The addition of zafirlukast to cetirizine improves the treatment of chronic urticaria in patients with positive autologous serum skin test results. *J Allergy Clin Immunol*, 113(1), 134-40. <https://doi.org/10.1016/j.jaci.2003.10.002>
- Baiardini, I., Braido, F., Bindslev-Jensen, C., Bousquet, P.J., Brzoza, Z., Canonica, G.W., ... Maurer, M. (2011). Recommendations for assessing patient-reported outcomes and health-related quality of life in patients with urticaria: a GA²LEN taskforce position paper. *Allergy*, 66, 840-844. <https://doi.org/10.1111/j.1398-9995.2011.02580.x>
- Banan, P., Butler, G., & Wu, J. (2014). Retrospective chart review in a cohort of patients with urticarial dermatitis. *Aust J Derm*, 55(2), 137-139. <https://doi.org/10.1111/ajd.12088>
- Baumgart, K., & Mullins, R. (2000). Use of hydroxychloroquine in refractory urticaria. *J Allergy Clin Immunol*, 105(1), S268-S269. [http://dx.doi.org/10.1016/S0091-6749\(00\)91223-5](http://dx.doi.org/10.1016/S0091-6749(00)91223-5)
- Beck, L.A., Marcotte, G.V., MacGlashan, D., Togias, A., & Saini, S. (2004). Omalizumab-induced reductions in mast cell Fc epsilon RI expression and function. *J Allergy Clin Immunol*, 114(3), 527-530. <https://doi.org/10.1016/j.jaci.2004.06.032>
- Ben-Shoshan, M., Blidnerman, I., & Raz, A. (2013). Psychosocial factors and chronic spontaneous urticaria: a systematic review. *Allergy*, 68(2), 31-41. <https://doi.org/10.1111/all.12068>
- Bernstein, J.A., Lang, D.M., Khan, D.A., Craig, T., Dreyfus, D., Hsieh, F., ... Wallace, D. (2014). The diagnosis and management of acute and chronic urticaria: 2014 update. *J Allergy Clin Immunol*, 133(5), 1270-1277. <https://doi.org/10.1016/j.jaci.2014.02.036>
- Bernstein, J.A., Kavati, A., Tharp, M.D., Ortiz, B., MacDonald, K., Denhaerynck, K. & Abraham, I.L. (2018). Effectiveness of omalizumab in adolescent and adult patients with chronic idiopathic/spontaneous urticaria: a systemic review of 'real-world' evidence. *Expert Opinion on Biological Therapy*, 18(4), 425-448. <https://doi.org/10.1080/14712598.2018.1438406>
- Berth-Jones, J., Hutchinson, P.E., Wicks, A.C., & Mitchell, V.E. (1988). Chronic urticaria with angio-oedema controlled by warfarin. *BMJ*. 297(6660), 1382-1383. <https://dx.doi.org/10.1136%2Fbmj.297.6660.1382>
- Bingham, C.O., Jones, P.E., Gober, L.M., Eckman, J.A., Sterba, P.M., & Saini SS. (2008). Targeted B-cell Depletion using Rituximab (Anti-CD20) in Recalcitrant Autoimmune Urticaria. *J Allergy Clin Immunol*, 121(2), S172. <https://doi.org/10.1016/j.jaci.2007.12.634>

Bleehan, S.S., Thomas, S.E., Greaves, M.W., Newton, J., Kennedy, C.T.C., Hindley, F., ... Howland, K. (1987). Cimetidine and chlorpheniramine in the treatment of chronic idiopathic urticaria: a multi-centre randomized double-blind study. *Br J Dermatol*, 117(1), 81-88.

<https://doi.org/10.1111/j.1365-2133.1987.tb04095.x>

Boonpiyathad, S., Fuengthong, R., & Sangasapaviriya, A. (2013). Effect of hydroxychloroquine treatment in the patients with anti-histamine refractory chronic spontaneous urticaria, randomised controlled trial. *Allergy*, 68(97), 62-62.

Boonpiyathad, T., & Sangasapaviliya, A. (2017). Hydroxychloroquine in the treatment of anti-histamine refractory chronic spontaneous urticaria, randomized single-blinded placebo-controlled trial and an open label comparison study. *Eur Ann Allergy Clin Immunol*, 49(5), 220-224.

<https://doi.org/10.23822/eurannaci.1764-1489.11>

Boubouka, C.D., Charissi, C., Kouimtzis, D., Kalogeromitros, D., Stavropoulos, P.G., & Katsarou, A. (2011). Treatment of autoimmune urticaria with low-dose cyclosporin A: A one-year follow-up. *Acta dermato-venereologica*, 91(1), 50-4. <https://doi.org/10.2340/00015555-0939>

Breneman, D.L. (1996). Cetirizine versus hydroxyzine and placebo in chronic idiopathic urticaria. *Ann Pharmacother*, 30(10), 1075-1079. <https://doi.org/10.1177%2F106002809603001001>

Broder, M.S., Raimundo, K., Antonova, E., & Chang, E. (2015). Resource Use and Costs in an Insured Population of Patients with Chronic Idiopathic/Spontaneous Urticaria. *Am J Clin Dermatol*, 16(4), 313-321. <https://doi.org/10.1007/s40257-015-0134-8>

Brown, N.A., & Carter, J.D. (2007). Urticular Vasculitis. *Curr Rheumatol*, 9(4), 312-319.
<https://doi.org/10.1007/s11926-007-0050-x>

Brüske, I., Standl, M., Weidinger, S., Klümper, C., Hoffmann, B., Schaaf, B., Herbarth I., ... Heinrich, J. (2013). Epidemiology of urticaria in infants and young children in Germany – Results from the German LISApplus and GINIplus Birth Cohort Studies. *Ped Allergy Immunol*. 25(1).
<https://doi.org/10.1111/pai.12146>

Bunselmeyer, B., Laubach, H.J., Schiller, M., Stanke, M., Luger, T.A., & Brehler, R. (2009). Incremental build-up food challenge - a new diagnostic approach to evaluate pseudoallergic reactions in chronic urticaria: a pilot study: stepwise food challenge in chronic urticaria. *Clin Exp Allergy*, 39(1)116-26. <https://doi.org/10.1111/j.1365-2222.2008.03110.x>

ASClA INFORMATION FOR HEALTH PROFESSIONALS

- Caffarelli, C., Cuomo B, Cardinale, F., Barberi, S., Dascola, C.P., Aqostinis, F., & Bernardini, R. (2013) Aetiological factors associated with chronic urticaria in children: a systematic review. *Acta Derm Venereol*, 93(3), 268-272. <https://doi.org/10.2340/00015555-1511>
- Cassano, N., D'Argento, V., Filotico, R., & Vena, G. (2005). Low dose dapsone in chronic idiopathic urticaria: Preliminary results of an open study. *Acta Derm Venereol*, 85(3), 254–255. <https://doi.org/10.1080/00015550510025560>
- Chakravarty, S.D., Yee, A.F., & Paget, S.A. (2011). Rituximab successfully treats refractory chronic autoimmune urticaria caused by IgE receptor autoantibodies. *J Allergy Clin Immunol*, 128(6), 1354-1355. <https://doi.org/10.1016/j.jaci.2011.08.023>
- Chansakulporn S, Pongpreuksa S, Sangacharoenkit P, Pacharn, P., Visitsunthorn, N. Vichyanond, Jirapongsananuruk, O. (2014). The natural history of chronic urticaria in childhood: a prospective study. *J Am Acad Dermatol*, 71(4), 663-668. <https://doi.org/10.1016/j.jaad.2014.05.069>
- Chua, S.L., & Gibbs, S. (2005). Chronic urticaria responding to subcutaneous heparin sodium. *Br J Dermatol*, 153(1), 216-217. <https://doi.org/10.1111/j.1365-2133.2005.06694.x>
- Church, M.K., Weller, C., Stock, P., & Maurer, M. (2011). Chronic spontaneous urticaria in children: Itching for insight. *Paediatr Allergy Immunol*, 22(1), 1-8. <https://doi.org/10.1111/j.1399-3038.2010.01120.x>
- Clarke, J.J., Secrest, A.M., Hull, C.M., Eliason, M.J. Leiferman, K.M., Gleich, G.J. & Powell, D.L. (2016). The effect of omalizumab dosing and frequency in chronic idiopathic urticaria: Retrospective chart review. *J Am Acad Dermatol*, 74(6), 1274-1276. <https://doi.org/10.1016/j.jaad.2015.12.052>
- Curto-Baredo, I., Spertino, J., Figueras-Nart, I., Expósito-Serrano, V., Guilabert, A., Melé-Ninot, G., ... Giménez-Arnau, A. (2018). Omalizumab updosing allows disease control in refractory patients with chronic spontaneous urticaria. *Br J Dermatol*, 179(1), 210-212. <https://doi.org/10.1111/bjd.16379>
- Di Lorenzo, G., Pacor, M.L., Mansueto, P., Pellitteri, E., Lo Bianco, C., Ditta, V., Martinelli, N., & Rini, B. (2004). Randomized placebo-controlled trial comparing desloratadine and montelukast in monotherapy and desloratadine plus montelukast in combined therapy for chronic idiopathic urticaria. *J Allergy Clin Immunol*, 114(3), 619-625. <https://doi.org/10.1016/j.jaci.2004.06.018>

Diaz-Angulo, S., Lopez-Hoyos, M., Munoz Casho P, Fernandez, M., Lopez-Escobar, M., Rodriguez, F., & Gonzalez-Lopez, M.A. (2016). Prevalence of thyroid autoimmunity in Spanish patients with chronic idiopathic urticaria: a case-control study involving 343 subjects. *J Eur Acad Dermatol Venereol*, 30, 692-693. <https://doi.org/10.1111/jdv.12979>

Ditto, A. M., (2007). Hydroxychloroquine in the treatment of 12 patients with severe chronic idiopathic urticaria; a case series. *J Allergy Clin Immunol*, 119(1).
<https://doi.org/10.1016/j.jaci.2006.12.147>

Doeglas, H.M., (1977). Dietary treatment of patients with chronic urticaria and intolerance to aspirin and food additives. *Dermatologica*, 54(5), 308-310.

Doshi, D.R., & Weinberger, M.M. (2009). Experience with cyclosporine in children with chronic idiopathic urticarial. *Pediatr Dermatol*, 26, 409-13. <https://doi.org/10.1111/j.1525-1470.2009.00869.x>

Duvall, L.A., Boackle, R.J., & King, R.G., (1986). Warfarin sodium therapy for chronic urticaria and angioedema. *South Med J*. 79(3), 389. <https://doi.org/10.1097/00007611-198603000-00039>

Eapen, A., & Kloepfer, K.M., (2018). Serum sickness-like reaction in a pediatric patient using omalizumab for chronic spontaneous urticaria. *Pedr All Immunol*, 29(4), 449-450.
<https://doi.org/10.1111/pai.12877>

Eastman, J., Wilde, N., & Jerath, M.R. (2014). Outcomes of Chronic Urticaria Patients Treated with Hydroxychloroquine. *J Allergy Clin Immunol*, 133(2), AB119.
[https://www.jacionline.org/article/S0091-6749\(13\)02345-2/pdf](https://www.jacionline.org/article/S0091-6749(13)02345-2/pdf)

Egeberg, A., Kofoed, K., Gislason, G.H., Vestergaard, C., & Thyssen, J.P. (2017). Cardiovascular risk is not increased in patients with chronic spontaneous urticaria: A retrospective population-based cohort study. *Acta Derm Venereol*, 97, 261-262.
https://www.medicaljournals.se/acta/content_files/files/pdf/97/2/4800.pdf

Ehlers, I., Niggemann, B., Binder, C., & Zuberbier, T. (1998). Role of nonallergic hypersensitivity reactions in children with chronic urticaria. *Allergy*, 53(11), 1074-1077.
<https://doi.org/10.1111/j.1398-9995.1998.tb03817.x>

Engin, B., & Ozdemir, M. (2008). Prospective randomised non-blinded clinical trial on the use of dapsone plus antihistamine vs. antihistamine in patients with chronic idiopathic urticaria. *JAD*, 22, 481–486. <https://doi.org/10.1111/j.1468-3083.2007.02510.x>

European Medicines Agency. Xolair summary of product characteristics. http://www.ema.europa.eu/docs/en_GB/document_library/EPAR_Product_Information/human/000606/WC500057298.pdf.

Erbagci, Z. (2002). The leukotriene receptor antagonist montelukast in the treatment of chronic idiopathic urticaria: a single-blind, placebo-controlled crossover clinical study. *J Allergy Clin Immunol*, 110, 484-488. <https://doi.org/10.1067/mai.2002.126676>

Fagiolo, U., Cancian, M., Bertollo, L., Peserico, A., & Amadori, A. (1999). Inhibitory effect of heparin on skin reactivity to autologous serum in chronic idiopathic urticaria. *J Allergy Clin Immunol*, 103(6), 1143-1147. [https://doi.org/10.1016/S0091-6749\(99\)70190-9](https://doi.org/10.1016/S0091-6749(99)70190-9)

Finn, A. F., Allen, P., Kaplan, M.D., Fretwell, R.F., Qu, R & Long, J. (1999). A double-blind, placebo-controlled trial of fexofenadine HCl in the treatment of chronic idiopathic urticaria. *J Allergy Clin Immunol*, 104(5), 1071-1078. [https://doi.org/10.1016/S0091-6749\(99\)70091-6](https://doi.org/10.1016/S0091-6749(99)70091-6)

Gach, J.E., Sabroe, R.A., Greaves, M.W., & Black, A.K. (2001). Methotrexate-responsive chronic idiopathic urticaria: a report of two cases. *Br J Dermatol*, 145, 340-343.
<https://doi.org/10.1046/j.1365-2133.2001.04330.x>

Gaig, P., Olona, M., Munoz Lejarazu, D., Caballero, M.T., Dominquez, F.J., Eschechipia, S., ... Ferrer, M. (2004). Epidemiology of urticaria in Spain. *J Investig Allergol Clin Immunol*, 14(3), 214-220. <https://pubmed.ncbi.nlm.nih.gov/15552715/>

Garg, G., & Thami, G.P. (2007). Comparative efficacy of cetirizine and levocetirizine in chronic idiopathic urticaria. *J Dermatolog Treat*, 18(1), 23-24. <https://doi.org/10.1186/1939-4551-7-33>

Godse, K.V. (2006). Oral montelukast monotherapy is ineffective in chronic idiopathic urticaria: a comparison with oral cetirizine. *Indian J Dermatol Venereol Leprol*, 72 (4), 312-314. <https://doi.org/10.4103/0378-6323.26735>

Goh, C.L., Wong, W.K., & Lim J. (1991). Cetirizine vs placebo in chronic idiopathic urticaria: a double blind randomized cross-over study. *Ann Acad Med Singapore*, 20(3), 328-30. <https://pubmed.ncbi.nlm.nih.gov/1681778/>

Goldsobel, A.B., Rohr, A.S., Siegel, S.C., Spector, S.L. Katz, R.M., Rachelefsky, G.S. Barr, R.J. (1986). Efficacy of doxepin in the treatment of chronic idiopathic urticaria. *J Allergy Clin Immunol*. 78(5), 867-873. [https://doi.org/10.1016/0091-6749\(86\)90232-0](https://doi.org/10.1016/0091-6749(86)90232-0)

Grant, J.A., Bernstein, D.I., Buckley, C.E., Chu, T., Fox, R.W., Rocklin, R.E. ... Karpenter, K.M. (1998). Double-blind comparison of terfenadine, chlorpheniramine and placebo in the treatment of

- chronic idiopathic urticaria. *J Allergy Clin Immunol*, 81(3), 574-579. [https://doi.org/10.1016/0091-6749\(86\)90232-0](https://doi.org/10.1016/0091-6749(86)90232-0)
- Grattan, C.E., Wallington, T.B., Warin, R.P., Kennedy, C.T.C., & Bradfield, J.W. (1986). A serological mediator in chronic idiopathic urticaria- a clinical, immunological and histological evaluation. *Br J Dermatol*, 114(5), 583-590. <https://doi.org/10.1111/j.1365-2133.1986.tb04065.x>
- Grattan, C.E., O'Donnell, B.F., Francis, D.M., Nimi, N., Barlow, R.J., Seed, P.T., ... Greaves, M.W. (2000). Randomized double blind study of cyclosporin in chronic 'idiopathic' urticaria. *Br J Dermatol*, 143(2), 365-372. <https://doi.org/10.1046/j.1365-2133.2000.03664.x>
- Grattan, C.E., Dawn, G., Gibbs, S., & Francis, D.M. (2003). Blood basophil numbers in chronic ordinary urticaria and healthy controls: diurnal variation, influence of loratadine and prednisolone and relationship to disease activity. *Clin Exp Allergy*, 33(3), 337-341. <https://doi.org/10.1046/j.1365-2222.2003.01589.x>
- Gray, P.E., Mehr, S., Katalaris, C.H., Wainstein, B.K., Star, A., Campbell, D., ... Ziegler, J.B. (2013). Salicylate elimination diets in children: is food restriction supported by the evidence? *Med J Aust*, 198(11), 600-602. <http://dx.doi.org/10.5694/mja12.11255>
- Greaves, M.W. (2000). Chronic urticaria in childhood. *Allergy*, 55(4), 309-320. <https://doi.org/10.1034/j.1398-9995.2000.00116.x>
- Green, J.P., & Maasyani, S. (1977). Tricyclic antidepressant drugs block histamine H₂ receptors in brain. (1977). *Nature*, 269(5624), 163-165. <https://doi.org/10.1038/269163a0>
- Groffik, A., Mitzel-Kaoukhov, H., Magerl, M., Maurer, M., & Staubach, P. (2011). Omalizumab-an effective and safe treatment of therapy-resistant chronic spontaneous urticaria. *Allergy*, 66(2), 303-305. <https://doi.org/10.1111/j.1398-9995.2010.02472.x>
- Guerra, L., Vincenzi, C., Marchesi, E., Tosti, A., Pretto, E., Bassi, R., ... De Costanza, F. (1994). Loratadine and cetirizine in the treatment of chronic urticaria. *J Eur Acad Dermatol Venereol*, 3(2), 148-152. <https://doi.org/10.1111/j.1468-3083.1994.tb00089.x>
- Guida, B., De Martino, C.D., De Martino, S.D., Tritto, G., Patella, V., Trio, R., ... D'Agostino L. (2000). Histamine plasma levels and elimination diet in chronic idiopathic urticaria. *Eur J Clin Nutr*, 54(2), 155-158. <https://doi.org/10.1038/sj.ejcn.1600911>
- Guillén-Aguinaga, S., Jáuregui Presa, I., Aguinaga-Ontoso, E., Guillén-Grima, F., & Ferrer, M. (2016). Updosing nonsedating antihistamines in patients with chronic spontaneous urticaria: a

- systematic review and meta-analysis. *Br J Dermatol* 175(6), 1153-1165.
<https://doi.org/10.1111/bjd.14768>
- Hale,T.W., (2012). Medications and Mothers' Milk. 15th Edition.
- Hall, A., & Marshall, G.D. (2013). A Retrospective review of hydroxychloroquine use in patients with chronic idiopathic urticaria (CIU). *Ann Allergy Asthma Immunol*, 111(5), A18-A19.
- Handa, S., Dogra, S., & Kumar, B. (2004). Comparative efficacy of cetirizine and fexofenadine in the treatment of chronic idiopathic urticaria. *J Dermatolog Treat*, 15(1), 55-57.
<https://doi.org/10.1080/09546630310013450>
- Hawro, T., Ohanyan, T., Schoepke, N., Metz, M., Peveling-Oberhag, A., Staubach, P., Maurer, M. & Weller, K. (2018). Comparison and Interpretability of the available Urticaria Activity Scores. *Allergy*, 73(1), 251-255. <https://doi.org/10.1111/all.13271>
- Henz, B.M., & Zuberbier, T. (1998). Most chronic urticaria is food-dependent, and not idiopathic. *Exp Dermatol*, 7(4), 139-142. <https://doi.org/10.1111/j.1600-0625.1998.tb00314.x>
- Hide, M., Francis, D.M., Grattan, C.E., Hakimi, J., Kochan, J.P. & Greaves, M.W. (1993). Autoantibodies against the high-affinity IgE receptor as a cause of histamine release in chronic urticaria. *N Engl J Med*, 328, 1599–1604. <https://doi.org/10.1056/nejm199306033282204>
- Holgate, S., Casale, T., Wenzel, S., Bousquet, J., Deniz, Y., & Reisner, C. (2005). The anti-inflammatory effects of omalizumab confirm the central role of IgE in allergic inflammation. *J Allergy Clin Immunol*, 115(3), 459-465. <https://doi.org/10.1016/j.jaci.2004.11.053>
- Hrabak, T., & Calabria, C.W. (2010). Multiple treatment cycles of high-dose intravenous immunoglobulin for chronic spontaneous urticaria. *Ann Allergy Asthma Immunol*, 105(3), 245. <https://doi.org/10.1016/j.anai.2010.05.018>
- Hsu, M.L., & Li, L.F. (2012). Prevalence of food avoidance and food allergy in Chinese patients with chronic urticaria. *Br J Dermatol*, 166(4), 747-752. <https://doi.org/10.1111/j.1365-2133.2011.10733.x>
- Iribarren, C., Tolstykh, I.V., Miller, M.K., Sobel, E., & Eisner, M.D. (2012). Adult asthma and risk of coronary heart disease, cerebrovascular disease, and heart failure: A prospective study of 2 matched cohorts. *Am J Epidemiol* 176(11), 1014-1024. <https://doi.org/10.1093/aje/kws181>
- Iribarren, C., Rahmaoui, A., Long, A.A., Szefler, S.J., Bradley, M.S., Carrigan, G., ... Rothman, K.J. (2017). Cardiovascular and cerebrovascular events among patients receiving omalizumab:

ASCIA INFORMATION FOR HEALTH PROFESSIONALS

Results from EXCELS, a prospective cohort study in moderate to severe asthma. *J Allergy Clin Immunol* 139(5), 1489-1495. <https://doi.org/10.1016/j.jaci.2016.07.038>

Jaffer, A.M. (1991). Sulfasalazine in the treatment of corticosteroid-dependent chronic idiopathic urticaria. *J Allergy Clin Immunol*, 88(6), 964-965. [https://doi.org/10.1016/0091-6749\(91\)90255-m](https://doi.org/10.1016/0091-6749(91)90255-m)

Jaffer, A.M., & Reid, R.T. (2002). Sulfasalazine in the treatment of chronic urticaria. *J Allergy Clin Immunol*, 109(1), S127. [https://doi.org/10.1016/0091-6749\(91\)90255-m](https://doi.org/10.1016/0091-6749(91)90255-m)

Jariwala, S.P., Moday, H., de Asis, M.L., Fodeman, J., Hudes, G., de Vos, G., & Rosenstreich, D. (2009). The Urticaria Severity Score: a sensitive questionnaire/index for monitoring response to therapy in patients with chronic urticarial. *Ann Allergy Asthma Immunol*, 102(6), 475-482. [https://doi.org/10.1016/s1081-1206\(10\)60120-2](https://doi.org/10.1016/s1081-1206(10)60120-2)

Jauregui, I., Ortiz de Frutos, F.J., Ferrer, M., Gimenez-Arnau, A., Sastre, J., Bartra, J., ... Valero, A. (2014). Assessment of severity and quality of life in chronic urticaria. *Journal of Investigational Allergology and Clin Immunol* 24(2), 80-86. <http://www.jiaci.org/issues/vol24issue2/2.pdf>

Jirapongsananuruk, O., Pongpreuksa, S., Sangacharoenkit, P., Visitsunthorn, N., & Vichyanond, P. (2010). Identification of the etiologies of chronic urticaria in children: a prospective study of 94 patients. *Pediatr Allergy Immunol*, 21(3), 508-514. <https://doi.org/10.1111/j.1399-3038.2009.00912.x>

Juhlin, L. (1981). Recurrent urticaria: clinical investigation of 330 patients. *Br J Dermatol*, 104(4), 369-381. <https://doi.org/10.1111/j.1365-2133.1981.tb15306.x>

Juhlin, L., & Arendt, C. (1988). Treatment of chronic urticaria with cetirizine dihydorchloride a non-sedating antihistamine. *Br J Dermatol*, 119(1), 67-72. <https://doi.org/10.1111/j.1365-2133.1988.tb07103.x>

Juhlin, L. (1991). Cetirizine in the treatment of chronic urticaria. *Clin Ther* 13 (1), 81-86. <https://pubmed.ncbi.nlm.nih.gov/1827614/>

Kalivas, J., Breneman, D., Tharp, M. Bruce, S., & Bigby, M. (1990). Urticaria: clinical efficacy of cetirizine in comparison with hydroxyzine and placebo. *J Allergy Clin Immunol* 86(2), 1014-1018. [https://doi.org/10.1016/s0091-6749\(05\)80246-5](https://doi.org/10.1016/s0091-6749(05)80246-5)

- Kameyoshi, Y., Tanaka, T., Mihara, S., Takahagi, S., Niimi, N., & Hide, M. (2007). Increasing the dose of cetirizine may lead to better control of chronic idiopathic urticaria: an open study of 21 patients. *Br J Dermatol*, 157(4), 803-804. <https://doi.org/10.1111/j.1365-2133.2007.08060.x>
- Kaplan, A.P. (2002). Clinical practice. Chronic urticaria and angioedema. *New England Journal of Medicine*, 346(3), 175-179. <https://doi.org/10.1056/nejmcp011186>
- Kaplan, A.P. (2004). Chronic urticaria: pathogenesis and treatment. *JACI* 114(3), 465-474. <https://doi.org/10.1016/j.jaci.2004.02.049>
- Kaplan, A.P., Spector, S.L., Meeves, S., Liao, Y., Verghese, S.t., & Georges, G. (2005). Once-daily fexofenadine treatment for chronic idiopathic urticaria: a multicenter, randomized, double-blind placebo-controlled study. *Ann Allergy Asthma Immunol*, 94(6), 662-669. [https://doi.org/10.1016/s1081-1206\(10\)61325-7](https://doi.org/10.1016/s1081-1206(10)61325-7)
- Kaplan, A.P., Joseph, K., Maykut, R.J., Geba, G.P., & Zeldin, R.K. (2008). Treatment of chronic autoimmune urticaria with omalizumab. *J Allergy Clin Immunol*, 122(3), 569-573. <https://doi.org/10.1016/j.jaci.2008.07.006>
- Kaplan, A.P. (2012). Treatment of chronic spontaneous urticaria. *Allergy Asthma Immunol Res*, 4(6), 326-331. <https://doi.org/10.4168/aaир.2012.4.6.326>
- Kaplan, A.P., Ferrer, M., Bernstein, J.A., Antonova, E., Trzaskoma, B., Raimundo, K., ... Zazzli, J.L. (2016). Timing and duration of omalizumab response in patients with chronic spontaneous urticaria. *J Allergy Clin Immunol*, 137(2), 474-481. <https://doi.org/10.1016/j.jaci.2015.08.023>
- Kapp, A., & Pichler, W.J. (2006). Levocetirizine is an effective treatment in patients suffering from chronic idiopathic urticaria: a randomized, double-blind, placebo-controlled, parallel, multicenter study. *Int J Dermatol* 45(4), 469-474. <https://doi.org/10.1111/j.1365-4632.2005.02609.x>
- Kawashima, M., & Harada, M. (2001). Efficacy and safety of fexofenadine HCl in Japanese patients with chronic idiopathic urticaria. *Int Arch Allergy Immunol*, 124, 343-345. <https://doi.org/10.1159/000053752>
- Kessel, A., Bamberger E., & Toubi, E. (2005). Tacrolimus in the treatment of severe chronic idiopathic urticaria: An open-label prospective study. *J Am Acad Dermatol*, 52(1), 145–148. <https://doi.org/10.1016/j.jaad.2004.09.023>
- Kessel, A., & Toubi, E. (2010). Cyclosporine-A in severe chronic urticaria: the option for long-term therapy. *Allergy*, 65(11), 1478-1482. <https://doi.org/10.1111/j.1398-9995.2010.02419.x>

ASClA INFORMATION FOR HEALTH PROFESSIONALS

- Khakoo, G., Sofianou-Katsoulis, A., Perkin, M.R., & Lack, G. (2008). Clinical features and natural history of physical urticaria in children. *Pediatr Allergy Immunol*, 19(4), 363-366.
<https://doi.org/10.1111/j.1399-3038.2007.00667.x>
- King, W., McCargar, L., Joneja, J.M., & Barr S.I. (2000). Benefits of a Histamine-Reducing Diet for Some Patients with Chronic Urticaria and Angioedema. *Can J Diet Pract Res*, 61(1), 24-26.
<https://pubmed.ncbi.nlm.nih.gov/11551340/>
- Kolkhir, P., Church, M.K., Weller, K., Metz, M., Schmetzer, O., & Maurer M. (2017). Autoimmune chronic spontaneous urticaria: What we know and what we do not know. *J Allergy Clin Immunol*, 139(6), 1772-1781. <https://doi.org/10.1016/j.jaci.2016.08.050>
- Konstantinou, G.N., Papadopoulos, N.G., Tavladaki, T., Tsekoura, T., Tsilimigaki, A., & Grattan, C.E. (2011). Childhood acute urticaria in northern and southern Europe shows a similar epidemiological pattern and significant meteorological influences. *Pediatr Allergy Immunol* 22, 36-42. <https://doi.org/10.1111/j.1399-3038.2010.01093.x>
- Konstantinou, G.N., Chioti, A.G., & Daniilidis, M. (2016). Self-reported hair loss in patients with chronic spontaneous urticaria treated with omalizumab: an under-reported, transient side effect? *Eur Ann Clin Immunol*, 48, 205-207. <http://www.eurannallergyimm.com/cont/journals-articles/445/volume-selfreported-hair-loss-patients-with-1188allasp1.pdf>
- Kosnik, M., (2011). Add-on montelukast in antihistamine-resistant chronic idiopathic urticaria. *Respir Med*, 105 (Suppl1), S84-S88. [https://doi.org/10.1016/s0954-6111\(11\)70018-x](https://doi.org/10.1016/s0954-6111(11)70018-x)
- Koti, I., Weller, K., Makris, M., Tiligada, E., Psaltopoulou, T., Papageorgiou, C., ... Maurer, M. (2013). Disease activity only moderately correlates with quality of life impairment in patients with chronic spontaneous urticaria. *Dermatology* 226(4), 371-379. <https://doi.org/10.1159/000351711>
- Kulthan, K., Gritiyarangsan, P., Sitakalin, C., Charuwichitratana, S., Korkij, W., Aunhacachoke, K., ... Kullavanijaya P. (2001). Multicenter study of the efficacy and safety of fexofenadine 60mg twice daily in 108 Thai patients with chronic idiopathic urticaria. *J Med Assoc Thai*, 84(2), 153-159. <https://pubmed.ncbi.nlm.nih.gov/11336071/>
- Kulthan, K., Chawekulrat, P., Komoltri, C., Hunnangkul, S., Tuchinda, P., Chularojanamontri, L., & Maurer, M. (2018). Cyclosporine for Chronic Spontaneous Urticaria: A Meta-Analysis and Systematic Review. *J Allergy Clin Immunol Pract*, 6(2), 586-599.
<https://doi.org/10.1016/j.jaip.2017.07.017>

- Lammert, J.K., & Robinson, D.M. (1996). Hydroxychloroquine (HCQ) in corticosteroid-dependent chronic urticaria (CDCU). *J Allergy Clin Immunol*, 97(1), 365.
- Ledford, D., Broder, M.S., Antonova, E., Omachi, T.A., Chang, E., & Luskin, A. (2016) Corticosteroid-related toxicity in patients with chronic idiopathic urticarial-chronic spontaneous urticaria. *Allergy Asthma Proc*, 37(6), 458-465. <https://doi.org/10.2500/aap.2016.37.3999>
- Lee, S.J., Ha, E.K., Jee, H.M., Lee, K.S., Lee, S.W., Kim, M.A., ... Han, H.Y., (2017). Prevalence and risk factors of urticaria with a focus on chronic urticaria in children. *Allergy Asthma Immunol Res*, 9(3), 212-219. <https://doi.org/10.4168/aair.2017.9.3.212>
- Legrain, V., Taieb , A., Sage, T., & Maleville, J. (1990). Urticaria in infants: a study of forty patients. *Pediatr Dermatol*, 7(2), 101-107. <https://doi.org/10.1111/j.1525-1470.1990.tb00663.x>
- Leznoff, A., & Sussman, G.L. (1989). Syndrome of idiopathic chronic urticaria and angioedema with thyroid autoimmunity: a study of 90 patients. *J Allergy Clin Immunol*, 84(1), 66-71. [https://doi.org/10.1016/0091-6749\(89\)90180-2](https://doi.org/10.1016/0091-6749(89)90180-2)
- Lieberman, P.L., Umetsu, D.T., Carrigan, G.J., & Rahmaoui, A. (2016). Anaphylactic reactions associated with omalizumab administration: Analysis of a case control study. *J Allergy Clin Immunol* 138(3), 913-915. <https://doi.org/10.1016/j.jaci.2016.03.030>
- Lindsay, K., Goulding, J., Solomon, M., & Broom, B. (2015). Treating chronic spontaneous urticaria using a brief 'whole person' treatment approach: a proof-of-concept study. *Clin Transl Allergy*, 5, 40. <https://ctajournal.biomedcentral.com/track/pdf/10.1186/s13601-015-0082-7.pdf>
- Liu, T.H., Lin, Y.R., Yang, K.C., Tsai, Y.G., Fu, Y.C., Wu, T.K., & Wu, H.P. (2010). Significant factors associated with severity and outcome of an initial episode of acute urticaria in children. *Pediatr Allergy Immunol*, 21(7), 1043-1051. <https://doi.org/10.1111/j.1399-3038.2010.01070.x>
- Long, A., Rahmaoui, A., Rothman, K.J., Guinan, E., Eisner, M., Bradley, M.S., ... Szeffler, S.J. (2014). Incidence of malignancy in patients with moderate-to-severe asthma treated with or without omalizumab. *J Allergy Clin Immunol*, 134(3), 560-567. <https://doi.org/10.1016/j.jaci.2014.02.007>
- Lynde, C., Sussman, G., Hébert, J., Gulliver, W., Yang, W.H., Chambenoit, O., ... Rihakova, L. (2017). Omalizumab Retreatment of Patients with Chronic Idiopathic Urticaria/Chronic Spontaneous Urticaria Following Return of Symptoms: Primary Results of the OPTIMA Study EADV. Poster abstract presented at the Clinical Dermatology Conference.

- Magen, E., & Mishal, J. (2012). The effect of L-thyroxine treatment on chronic idiopathic urticaria and autoimmune thyroiditis. *Int J Dermatol*, 51(1), 94-97. <https://doi.org/10.1111/j.1365-4632.2011.05172.x>
- Magerl, M., Pisarevskaja, D., Scheufele, R., Zuberbier, T., & Maurer, M. (2010). Effects of a pseudoallergen-free diet on chronic spontaneous urticaria: a prospective trial. *Allergy*, 65(1), 78-83. <https://doi.org/10.1111/j.1398-9995.2009.02130.x>
- Malanin, G., & Kalimo, K. (1989). Results of skin testing with food additives and the effect of an elimination diet in chronic and recurrent urticaria and recurrent angioedema. *Clin Exp Allergy*, 19(5), 539-543. <https://doi.org/10.1111/j.1365-2222.1989.tb02430.x>
- Mallipeddi, R., & Grattan, C.E. (2007). Lack of response of severe steroid-dependent chronic urticaria to rituximab. *Clin Exp Dermatol*, 32(3), 333-334. <https://doi.org/10.1111/j.1365-2230.2007.02365.x>
- McGirt, L.Y., Vasagar, K., Gober, L.M., Saini, S.S., & Beck, L.A. (2006). Successful treatment of recalcitrant chronic idiopathic urticaria with sulfasalazine. *Arch Dermatol*, 142(10), 1337-1342. <https://doi.org/10.1001/archderm.142.10.1337>
- Metz, M., Ohanyan, T., Church, M.K., & Maurer, M. (2014). Retreatment with omalizumab results in rapid remission in chronic spontaneous urticaria. *JAMA Dermatol*, 150, 288-290. <https://doi.org/10.1001/jamadermatol.2013.8705>
- Metz, M., Ohanyan, T., Church, M.K., & Maurer, M. (2014). Omalizumab is an effective and rapidly acting therapy in difficult-to-treat chronic urticaria: a retrospective clinical analysis. *J Dermatol Sci*, 73(1), 57-62. <https://doi.org/10.1016/j.jdermsci.2013.08.011>
- Mitzel-Kaoukhov, H., Staubach, P., & Muller-Brenne, T. (2010). Effect of high-dose intravenous immunoglobulin treatment in therapy-resistant chronic spontaneous urticaria. *Ann Allergy, Asthma & Immunol* 104, 253-258. <https://doi.org/10.1016/j.anai.2009.12.007>
- Mlynek, A., Zalewska-Janowska, A., Martus, P., Staubach, P., Zuberbier, T., Maurer, M. (2008). How to assess disease activity in patients with chronic urticaria? *Allergy* (6), 777-780. <http://dx.doi.org/10.1111/j.1398-9995.2008.01726.x>
- Monroe, E.W., Cohen, S.H., Kalbfleisch, J., & Schulz, C.I. (1981). Combined H1 and H2 antihistamine therapy in chronic urticaria. *Arch Dermatol*, 117(7), 404-407. <doi:10.1001/archderm.1981.01650070032018>

ASClA INFORMATION FOR HEALTH PROFESSIONALS

Monroe, E.W., Fox, R.W., Green, A.W., Izuno, C.G.T., Bernstein, D.I., Pleskow, W.W., Wills, I., & Brigante, J.R. (1988). Efficacy and safety of loratadine (10mg once daily) in the management of idiopathic chronic urticaria. *J Am Acad Dermatol*, 19(1), 138-139. [https://doi.org/10.1016/s0190-9622\(88\)80230-5](https://doi.org/10.1016/s0190-9622(88)80230-5)

Mathias, S.D., Crosby, R.D., Zazzali, J.L., Maurer, M., & Saini, S.S. (2012). Evaluating the minimally important difference of the urticaria activity score and other measures of disease activity in patients with chronic idiopathic urticaria. *Ann Allergy Asthma Immunol*, 108, 20-24. <https://doi.org/10.1016/j.anai.2011.09.008>

Maurer, M., Altrichter, S., Bieber, T., (2011). Efficacy and safety of omalizumab in patients with chronic urticaria who exhibit IgE against thyroperoxidase. *J Allergy Clin Immunol*, 128, 202e205–209e205. <https://doi.org/10.1016/j.jaci.2011.04.038>

Maurer, M., Rosen, K., Hsieh, H.J., Saini, S., Grattan, C., Gimenez-Arnau, A., ... Casale. T., (2013). Omalizumab for the treatment of chronic idiopathic or spontaneous urticaria. *N Engl J Med*; 368(10), 924-935. <https://doi.org/10.1056/nejmoa1215372>

Maurer, M., Church, M.K., & Weller, K. (2017). Chronic urticaria in children, still itching for insight. *JAMA Derm*, 153(12), 1221-1222. <http://doi:10.1001/jamadermatol.2017.3183>

Monroe, E.W. (1992). Relative efficacy and safety of loratadine, hydroxyzine, and placebo in chronic idiopathic urticaria and atopic dermatitis. *Clin Ther*, 14(1), 17-21. <https://pubmed.ncbi.nlm.nih.gov/1445478/>

Monroe, E.W. (1992). Relative efficacy and safety of loratadine, hydroxyzine and placebo in chronic idiopathic urticaria. *Arzneimittelforschung*, 42(9), 1119-1121. <https://pubmed.ncbi.nlm.nih.gov/1349509/>

Monroe, E., Finn, A., Patel, P., Guerrero, R., Ratner, P., & Bernstein, D. (2003). Efficacy and safety of desloratadine 5mg once daily in the treatment of chronic idiopathic urticaria: a double-blind, randomized, placebo-controlled trial. *J Am Acad Dermatol*, 48(4), 535-541. <https://doi.org/10.1067/mjd.2003.143>

Morgan, M., Cooke, A., Rogers, L., Adams-Huet, B., & Khan, D.A. (2014). Double-blind placebo-controlled trial of dapsone in antihistamine refractory chronic idiopathic urticaria. *J Allergy Clin Immunol Pract*, 2(5), 601-606. <https://doi.org/10.1016/j.jaip.2014.06.004>

- Mortureux, P., Leaute-Labreze, C., Legrain-Lifermann, V., Lamireau, T., Sarlangue, J., & Taieb, A. (1998). Acute urticaria in infancy and early childhood: a prospective study. *Arch Dermatol*, 134(3), 319-23. <https://doi.org/10.1001/archderm.134.3.319>
- Namazy, J., Cabana, M.D., Scheuerle, A.E., Thorp, J.M., Chen, H., Carrigan, G., Wang, Y., Veith, J., & Andrews, E.B. (2015). The Xolair pregnancy registry (EXPECT): The safety of omalizumab use during pregnancy. *J Allergy Clin Immunol*, 135(2), 407-412.
[https://www.jacionline.org/article/S0091-6749\(14\)01198-1/pdf](https://www.jacionline.org/article/S0091-6749(14)01198-1/pdf)
- Nelson, H.S., Raynolds, R., & Mason J. (2000). Fexofenadine HCl is safe and effective for treatment of chronic idiopathic urticaria. *Ann Allergy Asthma Immunol*, 84(5), 517-522.
[https://doi.org/10.1016/s1081-1206\(10\)62515-x](https://doi.org/10.1016/s1081-1206(10)62515-x)
- Netchiporuk, E., Sasseville, D., Moreau, L., Habel, Y., Rahme, E., & Ben-Shoshan, M. (2017). Evaluating comorbidities, natural history, and predictors of early resolution in a cohort of children with chronic urticaria. *JAMA Derm*, 153(12), 1236-1242.
<https://doi.org/10.1001/jamadermatol.2017.3182>
- Nettis, E., Dambra, P., D'Oronzio, L., Loria, M.P., Ferrannini, A., & Tursi, A. (2001). Comparison of montelukast and fexofenadine for chronic idiopathic urticaria. *Arch Dermatol*, 137(1) 99-100.
- Nettis, E., Pannofino, A., Cavallo, E., Ferrannini, A., & Tursi, A. (2003). Efficacy of montelukast, in combination with loratadine, in the treatment of delayed pressure urticaria. *J Allergy Clin Immunol*, 112(1), 212-213. <https://doi.org/10.1067/mai.2003.1559>
- Nettis, E., Colanardi, M.C., Paradiso, M.T. & Ferrannini, A. (2004). Desloratadine in combination with montelukast in the treatment of chronic urticaria: a randomized, double-blind, placebo-controlled study. *Clin Exp Allergy*, 34(9), 1401-1407. <https://doi.org/10.1111/j.1365-2222.2004.02019.x>
- Nettis, E., Colanardi, M.C., Barra, L., Ferrannini, A., Vacca, A., & Tursi, A. (2006). Levocetirizine in the treatment of chronic idiopathic urticaria: a randomized, double-blind, placebo-controlled study. *Br J Dermatol*, 154(3), 533-538. <https://doi.org/10.1111/j.1365-2133.2005.07049.x>
- Noshela Ghazanfar, M., & Thomsen, S.F. (2017). Transient hair loss in patients with chronic spontaneous urticaria treated with omalizumab. *Eur Ann Allergy Clin Immunol*, 49(6), 284-285.
<https://doi.org/10.23822/eurannaci.1764-1489.08>
- O'Donnell, B.F., Barr, R.M., Black, A.K., Francis, D.M., Kermani, F., Niimi, N., ... Greaves, M.W. (1998). Intravenous immunoglobulin in autoimmune chronic urticaria. *Br J Dermatol* 138(1), 101-106.
[Intravenous immunoglobulin in autoimmune chronic urticaria](#)

- Ohanyan, T., Schoepke, N., Bolukbasi, B., Metz, M., Hawro, T., Zuberbier, T., ... Weller, K. (2017). Responsiveness and minimal important difference of the urticaria control test. *J Allergy Clin Immunol*, 140(6), 1710-1713. <https://doi.org/10.1016/j.jaci.2017.04.050>
- Orden, R.A., Timble, H., & Saini, S.S. (2014). Efficacy and safety of sulfasalazine in patients with chronic idiopathic urticaria. *Ann Allergy Asthma Immunol*. 112(1), 64-70. <https://doi.org/10.1016/j.anai.2013.09.028>
- Ortonne, J.P., Grob, J.J., Auquier, P., & Dreyfus, I. (2007). Efficacy and safety of desloratadine 5mg once daily in the treatment of chronic idiopathic urticaria: a double-blind, placebo-controlled, multicenter trial. *Am J Clin Dermatol*, 8(1), 37-42. <https://doi.org/10.1067/mjd.2003.143>
- Pacor, M.L., Di Lorenzo, G., & Corrocher, R. (2001). Efficacy of leukotriene receptor antagonist in chronic urticaria. A double-blind, placebo-controlled comparison of treatment with montelukast and cetirizine in patients with chronic urticaria with intolerance to food additive and/or acetylsalicylic acid. *Clin Exp Allergy*, 31(10), 1607-1614. <https://doi.org/10.1046/j.1365-2222.2001.01189.x>
- Parslew, R., Pryce D, Ashworth J, & Friedmann PS. (2000). Warfarin treatment of chronic idiopathic urticaria and angio-oedema. *Clin Exp Allergy*, 30(8), 1161-1165. <https://doi.org/10.1046/j.1365-2222.2000.00857.x>
- Paul, E., & Bodeker, R.H. (1986). Treatment of chronic urticaria with terfenadine and ranitidine. A randomized double-blind study in 45 patients. *Eur J Clin Pharmacol*, 31(3), 277-280. <https://doi.org/10.1007/bf00981123>
- Perez, A., Woods, A., & Grattan, C.E. (2010). Methotrexate: a useful steroid-sparing agent in recalcitrant chronic urticaria. *Br J Dermatol*, 162(1), 191-194. <https://doi.org/10.1111/j.1365-2133.2009.09538.x>
- Pitt, T., Warrington, R., & Kalicinsky, C. (2010). Adjunctive treatment of chronic idiopathic urticaria and angioedema with sulfasalazine. *Allergy Asthma Clin Immunol*, 6(1), 25. <https://aacijournal.biomedcentral.com/track/pdf/10.1186/1710-1492-6-S1-P25.pdf>
- Pongonis, R.M., & Fahrenholz, J.M. (2012). Efficacy of Hydroxychloroquine in the Treatment of 19 Patients with Antihistamine-refractory Chronic Urticaria. *J Allergy Clin Immunol*, 129(2), AB224. <https://doi.org/10.1016/j.jaci.2011.12.122>

ASClA INFORMATION FOR HEALTH PROFESSIONALS

- Potter, P.C., Kapp, A., Maurer, M., Guillet, G., Jian, A.M., Hauptmann, P., & Finlay, A.Y. (2009). Comparison of the efficacy of levocetirizine 5mg and desloratadine 5mg in chronic idiopathic urticaria patients. *Allergy*, 64(4), 596-604. <https://doi.org/10.1111/j.1398-9995.2008.01893.x>
- Powell, R.J., Du Toit, G.L., Siddique, N., Leech, S.C., Dixon, T.A., Clark, A.T., ... Huber, P.A. (2007). BSACI guidelines for the management of chronic urticaria and angio-oedema. *Clin Exp Allergy*, 37(5), 631-650. <https://doi.org/10.1111/cea.12494>
- Puccetti, A., Bason, C., Simeoni, S., Millo, E., Tinazzi, E., Beri, R., ... Lundari C. (2005). In chronic idiopathic urticaria autoantibodies against Fc epsilonRII/CD23 induce histamine release via eosinophil activation. *Clin Exp Allergy*, 35(12), 1599-1607. <https://doi.org/10.1111/j.1365-2222.2005.02380.x>
- Raghavendran, R.R., Humphreys, F., & Kaur, M.R. (2014). Successful use of mycophenolate mofetil to treat severe chronic urticaria in a patient intolerant to cyclosporin. *Clin Exp Dermatol*, 39(1), 68-69. <https://doi.org/10.1111/ced.12156>
- Rajan, J., Simon, R., & Bosso, J. (2014). Prevalence of sensitivity to food and drug additives in patients with chronic idiopathic urticaria. *J Allergy Clin Immunol Pract*, 2(2), 168-171. <https://doi.org/10.1016/j.jaip.2013.10.002>
- Reese, I., Zuberbier, T., Bunselmeyer, B., Erdmann, S., Henzgen, M., Fuchs, T., ... Werfel T. J. (2009). Diagnostic approach for suspected pseudoallergic reaction to food ingredients. *Dtsch Dermatol Ges*, 7(1), 70-77. <https://doi.org/10.1111/j.1610-0387.2008.06894.x>
- Reeves, G., Boyle, M., Bonfield, J., Dobson, P., & Loewenthal, M. (2004). Impact of hydroxychloroquine therapy on chronic urticaria: chronic autoimmune urticaria study and evaluation. *Internal Medicine Journal*, 34(4), 182-186. <https://doi.org/10.1111/j.1444-0903.2004.00532.x>
- Reider, N., & Egger, C. (2009). Infliximab in the treatment of refractory chronic urticaria. *J Am Acad Dermatol*, 60(3), AB39. <https://doi.org/10.1016/j.jaad.2008.11.193>
- Reimers, A., Pichler, C., Helbling, A., Pichler, W.J., & Yawalkar, N. (2002). Zafirlukast has no beneficial effects in the treatment of chronic urticaria. *Clin Exp Allergy*, 32(12), 1763-1768 <https://doi.org/10.1046/j.1365-2222.2002.01536.x>
- Richelson, E. (1979). Tricyclic antidepressants and histamine H1 receptors. *Mayo Clin Proc*, 54, 669. <https://pubmed.ncbi.nlm.nih.gov/39202/>

- Ring, J., Hein, R., Gauger, A., Bronsky, E., & Miller, B. (2001). Once-daily desloratadine improves the signs and symptoms of chronic idiopathic urticaria: a randomized, double-blind placebo-controlled study. *Int J Dermatol*, 40(1), 72-76. <https://doi.org/10.1046/j.1365-4362.2001.00186.x>
- Rorie, A., Goldner, W.S., Lyden, E., & Poole, J.A. (2014). Beneficial role for supplemental vitamin D3 treatment in chronic urticaria: a randomized study. *Ann Allergy Asthma Immunol*, 112(4), 376-382. <https://doi.org/10.1016/j.anai.2014.01.010>
- Rudzki, E., Czubalski, K., & Grzywa, Z. (1980). Detection of urticaria with food additives intolerance by means of diet. *Dermatologica*, 161(1), 57-62. <https://doi.org/10.1159/000250331>
- Rumbyrt, J.S., Katz, J.L., & Schocket AL. (1995). Resolution of chronic urticaria in patients with thyroid autoimmunity. *J Allergy Clin Immunol*. 96(6 Pt 1), 901-905. [https://doi.org/10.1016/S0091-6749\(95\)70226-1](https://doi.org/10.1016/S0091-6749(95)70226-1)
- Sabroe, R.A., & Greaves, M.W. (2006). Chronic idiopathic urticaria with functional autoantibodies: 12 years on. *Br J Dermatol*. 154(5), 813-819. <https://doi.org/10.1111/j.1365-2133.2006.07183.x>
- Sackesen, C., Sekerel, B.E., Orhan, F., Kocabas, C.N., Tuncer, A., & Adalioglu, G. (2004). The etiology of different forms of urticaria in childhood. *Pediatr Dermatol*, 21(2), 102-108. <https://doi.org/10.1111/j.0736-8046.2004.21202.x>
- Sagi, L., Solomon, M., Baum, S., Lyakhovitsky, A., Trau, H., & Barzilai, A. (2011). Evidence for methotrexate as a useful treatment for steroid-dependent chronic urticaria. *Acta Dermato Venereol*, 91(3), 303-306. <https://doi.org/10.2340/00015555-1080>
- Sahiner, U.M., Civelek, E., Tuncer, A., Yavuz, S.T., Karabulut, E., Sackesen, C., & Sekerel, B.E. (2011). Chronic urticaria: etiology and natural course in children. *Int Arch Allergy Immunol*, 156(2), 224-230. <https://doi.org/10.1159/000322349>
- Saini, S.S., & MacGlashan, D. (2002). How IgE upregulates the allergic response. *Curr Opin Immunol*, 14(6), 694-697. [https://doi.org/10.1016/s0952-7915\(02\)00404-1](https://doi.org/10.1016/s0952-7915(02)00404-1)
- Saini, S. S., Rosen, K.E., Hsieh, H.J., Wong, D.A., Conner, E., Kaplan, A., Spector, S., & Maurer, M. (2011). A randomized, placebo-controlled, dose-ranging study of single-dose omalizumab in patients with H1-antihistamine-refractory chronic idiopathic urticaria. *J Allergy Clin Immunol*, 128(3), 567-573. <https://doi.org/10.1016/j.jaci.2011.06.010>
- Saini, S.S., Bindslev-Jensen, C., Maurer, M., Grob, J.J., Bulbul Baskan, E., Bradley, M.S., ... Rosen, K. (2015). Efficacy and safety of omalizumab in patients with chronic idiopathic/spontaneous

- urticaria who remain symptomatic on H1 antihistamines: a randomized, placebo-controlled study. *J Invest Dermatol*, 135(1), 67-75. <https://dx.doi.org/10.1038%2Fjid.2014.306>
- Samarasinghe, V., & Marsland, A.M. (2012). Class action of oral coumarins in the treatment of a patient with chronic spontaneous urticaria and delayed-pressure urticaria. *Clin Exp Dermatol*, 37(7), 741-743. <https://doi.org/10.1111/j.1365-2230.2011.04281.x>
- Sand, F.L., & Thomsen, S.F. (2013). TNF-Alpha Inhibitors for Chronic Urticaria: Experience in 20 Patients. *J Allergy (Cairo)*, 130905. <https://dx.doi.org/10.1155%2F2013%2F130905>
- Shahar, E., Bergman, R., Guttman-Yassky, E., & Pollack, S. (2006). Treatment of severe chronic idiopathic urticaria with oral mycophenolate mofetil in patients not responding to antihistamines and/or corticosteroids. *International Journal of Dermatology*, 45, 1224-1227. <https://doi.org/10.1111/j.1365-4632.2006.02655.x>
- Shamsi, Z., & Hindmarch, I. (2000). Sedation and antihistamines: a review of inter-drug differences using proportional impairment ratios. *Human Psychopharmacol*, 15(s1), s3-s30. [https://doi.org/10.1002/1099-1077\(200010\)15:1+%3C::aid-hup247%3E3.0.co;2-s](https://doi.org/10.1002/1099-1077(200010)15:1+%3C::aid-hup247%3E3.0.co;2-s)
- Shamsi, Z., & Hindmarch, I. (2008). Sedation and antihistamines: a review of inter-drug differences using proportional impairment ratios. *Human Psychopharmacol*, 23(7), 555-570. <https://doi.org/10.1002/hup.962>
- Sharma, V.K., Singh, S., Ramam, M., Kumawat, M., & Kumar, R. (2014). A randomized placebo-controlled double-blind pilot study of methotrexate in the treatment of H1 antihistamine-resistant chronic spontaneous urticaria. *Indian J Dermatol Venereol Leprol*, 80(2), 122-128. <https://doi.org/10.4103/0378-6323.129382>
- Sharpe, G.R., & Shuster, R. (1993). In dermatographic urticaria H2 receptor antagonists have a small but therapeutically irrelevant additional effect compared with H1 antagonists alone. *Br J Dermatol*, 129(5), 575-579. <https://doi.org/10.1111/j.1365-2133.1993.tb00487.x>
- Shin, M., & Lee, S., (2017). Prevalence and causes of childhood urticaria. *Allergy, Asthma & Immunology Research*, 9(3), 189-190. <https://doi.org/10.4168/aair.2017.9.3.212>
- Siebenhaar, F., Degener, F., Zuberbier, T., Martus, P. & Maurer, M. (2009). High-dose desloratadine decreases wheal volume and improves cold provocation thresholds compared with standard dose treatment in patients with acquired cold urticaria: a randomized, placebo-controlled crossover study. *J Allergy Clin Immunol*, 123(3), 672-679. <https://doi.org/10.1016/j.jaci.2008.12.008>

- Simons, F.E. (2001). Prevention of acute urticaria in young children with atopic dermatitis. *J Allergy Clin Immunol*, 107(4), 703-706. <https://doi.org/10.1067/mai.2001.113866>
- Spector, S.L., Shikiar, R., Harding, G., Meeves, S., & Leahy, M.J. (2007). The effect of fexofenadine hydrochloride on productivity and quality of life in patients with chronic idiopathic urticaria. *Cutis*, 79(2), 157-162. <https://europepmc.org/article/med/17388220>
- Staevska, M., Popov, T.A. Kralimarkova, T., Lazatova, C., Kraeva, S., Popova, D., ... Church, M.K. (2010). The effectiveness of levocetirizine and desloratadine in up to 4 times conventional doses in difficult-to-treat urticaria. *J Allergy Clin Immunol*, 125(3), 676-682. <https://doi.org/10.1016/j.jaci.2009.11.047>
- Staevska, M., Gugutkova, M., Lazarova, C. Kralimarkova, T., Dimitrov, V., Zuberbier, T., Church, M.K. & Popov, T.A. (2014). Night-time sedating H1-antihistamine increases daytime somnolence but not treatment efficacy in chronic spontaneous urticaria: a randomized controlled trial. *Br J Dermatol*, 171(1), 148-154. <https://doi.org/10.1111/bjd.12846>
- Staubach, P., Dechene, M., Metz, M., Magerl, M., Siebenhaar, F., Weller, K., ... Maurer, M. (2011). High prevalence of mental disorders and emotional distress in patients with chronic spontaneous urticaria. *Acta Derm Venereol*, 91(5), 557-561. <https://doi.org/10.2340/00015555-1109>
- Staubach, P., Metz, M., Chapman-Rothe, N., Sieder, C., Brautigam, M., Canvin, J., & Maurer, M. (2016). H1-antihistamine-resistant chronic spontaneous urticaria patients: results from X-ACT, a randomized controlled trial. *Allergy*, 71(8), 1134-1144. <https://doi.org/10.1111/all.12870>
- Sussman, G., Hébert, J., Barron, C., Bian, J., Caron-Guay, R.M., Laflamme, S., & Stern, S. (2014). Real-life experiences with omalizumab for the treatment of chronic urticaria. *Ann Allergy Asthma Immunol*, 112(2), 170-174. <https://doi.org/10.1016/j.anai.2013.12.005>
- Tal, Y., Toker, O., Agmon-Levin, N., & Shalit, M. (2014). Azathioprine as a therapeutic alternative for refractory chronic urticaria. *Int J Dermatol*, 54(3), 367-369. <http://dx.doi.org/10.1111/ijd.12536>
- Tedeschi, A. (2009). Paradoxical exacerbation of chronic urticaria by H1-antihistamines and montelukast. *Europ Ann Allergy Clin Immunol*, 41(6), 187-189. <http://www.eurannallergyimm.com/cont/journals-articles/175/volume-paradoxical-exacerbation-chronic-urticaria-hantihistamines-470allaspl.pdf>
- Thompson, A.K., Finn, A.F., & Schoenwetter, W.F. (2000). Effect of 60mg twice-daily fexofenadine HCl on quality of life work and classroom productivity, and regular activity in patients with chronic

idiopathic urticaria. *J Am Acad Dermatol* 43(1 Pt 1), 24-30.

<https://doi.org/10.1067/mjd.2000.104680>

Tondury, B., Muehleisen, B., Ballmer-Weber, B.K., Hofbauer, G., Schmid-Grendelmeier, P., French, L., & Buchi, S. (2011). The Pictorial Representation of Illness and Self Measure (PRISM) instrument reveals a high burden of suffering in patients with chronic urticaria. *J Investig Allergol Clin Immunol*, 21(2), 93-100. <http://dx.doi.org/10.5167/uzh-48861>

Toubi E, Kessel A, Avshovich N, Bamber E, Sabo E, Nusem D, Panasoff J. (2004). Clinical and laboratory parameters in predicting chronic urticaria duration: a prospective study of 139 patients. *Allergy*, 59, 869-873. <https://doi.org/10.1111/j.1398-9995.2004.00473.x>

Vadasz, Z., Tal, Y., Rotem, M., Shichter-Confino, V., Mahlab-Guri, K. Graif, Y., ... Sthoeger, Z. (2017). Omalizumab for severe chronic spontaneous urticaria: Real-life experiences of 280 patients. *J Allergy Clin Immunol Pract*, 5(6), 1743-1745. <https://doi.org/10.1016/j.jaip.2017.08.035>

Van den Elzen, M., van Os-Medendorp, H., van den Brink, I., van den Hurk, K., Kouznetsova, O.I., ... Knulst, A.C. (2017). Effectiveness and safety of antihistamines up to fourfold or higher in treatment of chronic spontaneous urticaria. *Clinical and Translational Allergy*, 7(4). <https://dx.doi.org/10.1186%2Fs13601-017-0141-3>

Varghese, R., Rajappa, M., Chandrashekhar, L., Kattimani, S., Archana, M., Munisamy, M., Revathy, G., & Thappa D. M. (2016) Association among stress, hypocortisolism, systemic inflammation, and disease severity in chronic urticaria. *Ann Allergy Asthma Immunol*. 116(4), 344-348. <https://doi.org/10.1016/j.anai.2016.01.016>

Vena, G., Cassano, N., Colombo, D., Peruzzi, E., & Pigatto, P. (2006). Cyclosporine in chronic idiopathic urticaria: a double-blind, randomised, placebo-controlled trial. *J Am Acad Dermatol* 55, 705–709. <https://doi.org/10.1016/j.jaad.2006.04.078>

Verschave, A., Stevens, E., & Degreef, H. (1983). Pseudo-allergen-free diet in chronic urticaria. *Dermatologica*, 167(5), 256-259. <https://doi.org/10.1159/000249792>

Wagner, N., Dirk, D., Peveling-Oberhag, A., Reese, I., Rady-Pizarro, U., Mitzel, H., & Staubach, P. (2017). A Popular myth - low-histamine diet improves chronic spontaneous urticaria - fact or fiction? *J Eur Acad Dermatol Venereol*, 31(4), 650-655. <https://doi.org/10.1111/jdv.13966>

Wan, K.S. (2009). Efficacy of leukotriene receptor antagonist with an anti-H1 receptor antagonist for treatment of chronic idiopathic urticaria. *J Dermatolog Treat*, 20, 194-197.

<https://doi.org/10.1080/09546630802607495>

- Wan, K.S., & Chang, (2014). Efficacy of leukotriene receptor antagonist with anti-H1 receptor antagonist plus anti-H2 receptor antagonist for treatment of refractory chronic idiopathic urticaria. *J Dermatolog Treat*, 25(6), 459-461. <https://doi.org/10.3109/09546634.2013.849791>
- Weller, K.G., Magerl, M., Tohme, N., Martus, P., Krause, K., Metz, M., Staubach, P., & Maurer M. (2013). Development, Validation and Initial Results of the Angioedema Activity Score. *Allergy*, 68(9), 1185-1192. <https://doi.org/10.1111/all.12209>
- Weller, K., Groffik, A., Church, M.K., Hawro, T., Krause, K., Metz, M., ... Maurer, M. (2014). Development and validation of the Urticaria Control Test: a patient-reported outcome instrument for assessing urticaria control. *J Allergy and Clin Immunol*, 133(5), 1365-1372. <https://doi.org/10.1016/j.jaci.2013.12.1076>
- Wetter, D.A., Davis, M.D., Yiannias, J.A., Gibson, L.E., Dahl, M.V., el-Azhary, R.A., ... Pittelkow, M.R. (2005). Effectiveness of intravenous immunoglobulin therapy for skin disease other than toxic epidermal necrolysis: a retrospective review of Mayo Clinic experience. *Mayo Clinic Proceedings* 80, 41-47. <https://doi.org/10.4065/80.1.41>
- Wilson, L.H., Eliason, M.J., Leiferman, K.M., Hull, C.M., & Powell, D.L. (2011). Treatment of refractory chronic urticaria with tumor necrosis factor alfa inhibitors. *J Am Acad Dermatol*. 64(6), 1221-1222. <https://doi.org/10.1016/j.jaad.2009.10.043>
- Zimmerman, A.B., Berger, E.M., Elmariyah, S.B., & Soter, N.A. (2012). The use of mycophenolate mofetil for the treatment of autoimmune and chronic idiopathic urticaria: experience in 19 patients. *Journal of the American Academy of Dermatology*, 66, 767-770. <https://doi.org/10.1016/j.jaad.2011.06.004>
- Zitelli, K.B., & Cordoro, K.M. (2011). Evidence-based evaluation and management of chronic urticaria in children. *Pediatr Dermatol*, 28(6), 629-639. <https://doi.org/10.1111/j.1525-1470.2011.01518.x>
- Zuberbier, T., Chantraine-Hess, S., Hartmann, K., & Czarnetzki, B.M. (1995). Pseudoallergen-free diet in the treatment of chronic urticaria. A prospective study. *Acta Derm Venereol*. 75(6), 484-487. <https://doi.org/10.2340/000155575484487>
- Zuberbier, T., Munzberger, C., Haustein, U., Trippas, E., Burtin, B., Mariz, S.D. & Henz, B.M. (1996). Double-blind crossover study of high-dose cetirizine in cholinergic urticaria. *Dermatology*, 193(4), 324-327. <https://doi.org/10.1159/000246281>
- Zuberbier, T. (2001). The role of allergens and pseudoallergens in urticaria. *J Investig Dermatol Symp Proc*, 6(2), 132-134. <https://doi.org/10.1046/j.0022-202x.2001.00024.x>

Zuberbier, T., Asero, R., Bindslev-Jensen, G., Canonica, G., Church, M.K., Gimenez-Arnau, A., ...

Maurer, M. (2009). The EAACI/GA²LEN/EDF/WAO guideline: definition, classification and diagnosis of urticaria. *Allergy*, 64(10), 1417-1426. <https://doi.org/10.1111/j.1398-9995.2009.02179.x>

Zuberbier, T., Balke, M., Worm, M., Edenharder, G., & Maurer, M. (2010). Epidemiology of Urticaria; a representative cross-sectional population survey. *Clin Exp Dermatol*, 35(8), 869-783.

<https://doi.org/10.1111/j.1365-2230.2010.03840.x>

Zuberbier, T., Aberer, W., Asero, R., Abdul Latiff, A.H., Baker, D., Ballmer-Weber, B., ... Maurer, M. (2018). The EAACI/GA²LEN/EDF/WAO Guideline for the Definition, Classification, Diagnosis and Management of Urticaria. The 2017 Revision and Update. *Allergy* 73(7), 1393-1414.

<https://doi.org/10.1111/all.13397>